

Cinnamon may be used to halt progression of Parkinson's disease, study suggests

Date: July 9, 2014

Source: Rush University Medical Center

Summary: Using cinnamon, a common food spice and flavoring material, can reverse the biomechanical, cellular and anatomical changes that occur in the brains of mice with Parkinson's disease (PD), neurological scientists have found. "This could potentially be one of the safest approaches to halt disease progression in Parkinson's patients," the study's lead researcher said.

FULL STORY

Neurological scientists at Rush University Medical Center have found that using cinnamon, a common food spice and flavoring material, can reverse the biomechanical, cellular and anatomical changes that occur in the brains of mice with Parkinson's disease (PD). The results of the study were recently published in the June 20 issue of the *Journal of Neuroimmune Pharmacology*.

"Cinnamon has been used widely as a spice throughout the world for centuries," said Kalipada Pahan, PhD, study lead researcher and the Floyd A. Davis professor of neurology at Rush. "This could potentially be one of the safest approaches to halt disease progression in Parkinson's patients."

"Cinnamon is metabolized in the liver to sodium benzoate, which is an FDA-approved drug used in the treatment for hepatic metabolic defects associated with hyperammonemia," said Pahan. It is also widely used as a food preservative due to its microbiocidal effect.

Chinese cinnamon (*Cinnamomum cassia*) and original Ceylon cinnamon (*Cinnamomum verum*) are two major types of cinnamon that are available in the US.

"Although both types of cinnamon are metabolized into sodium benzoate, by mass spectrometric analysis, we have seen that Ceylon cinnamon is much more pure than Chinese cinnamon as the latter contains coumarin, a hepatotoxic molecule," said Pahan.

"Understanding how the disease works is important to developing effective drugs that protect the brain and stop the progression of PD," said Pahan. "It is known that some important proteins like Parkin and DJ-1 decrease in the brain of PD patients."

The study found that after oral feeding, ground cinnamon is metabolized into sodium benzoate, which then enters into the brain, stops the loss of Parkin and DJ-1, protects neurons, normalizes neurotransmitter levels, and improves motor functions in mice with PD.

This research was supported by grants from National Institutes of Health.

"Now we need to translate this finding to the clinic and test ground cinnamon in patients with PD. If these results are replicated in PD patients, it would be a remarkable advance in the treatment of this devastating neurodegenerative disease," said Dr. Pahan.

Parkinson's disease is a slowly progressive disease that affects a small area of cells within the mid-brain known as the substantia nigra. Gradual degeneration of these cells causes a reduction in a vital chemical neurotransmitter, dopamine. The decrease in dopamine results in one or more of the classic signs of Parkinson's disease that includes: resting tremor on one side of the body; generalized slowness of movement; stiffness of limbs; and gait or balance problems. The cause of the disease is unknown. Both environmental and genetic causes of the disease have been postulated.

Parkinson's disease affects about 1.2 million patients in the United States and Canada. Although 15 percent of patients are diagnosed before age 50, it is generally considered a disease that targets older adults, affecting one of every 100 persons over the age of 60. This disease appears to be slightly more common in men than women.

Story Source:

Materials provided by **Rush University Medical Center**. *Note: Content may be edited for style and length.*

Journal Reference:

1. Saurabh Khasnavis, Kalipada Pahan. **Cinnamon Treatment Upregulates Neuroprotective Proteins Parkin and DJ-1 and Protects Dopaminergic Neurons in a Mouse Model of Parkinson's Disease.** *Journal of Neuroimmune Pharmacology*, 2014; DOI: 10.1007/s11481-014-9552-2

Cite This Page:


MLA

APA

Chicago

Rush University Medical Center. "Cinnamon may be used to halt progression of Parkinson's disease, study suggests." ScienceDaily. ScienceDaily, 9 July 2014.
<www.sciencedaily.com/releases/2014/07/140709095257.htm>.

RELATED STORIES



Gut Microbes Promote Motor Deficits in a Mouse Model of Parkinson's Disease

Dec. 1, 2016 — Gut microbes may play a critical role in the development of Parkinson's-like movement disorders in genetically predisposed mice, researchers report. Antibiotic treatment reduced motor deficits ... **read more »**

New Gene Involved in Parkinson's Disease Found, Finding That May Result in New Treatments

June 4, 2014 — A new gene involved in Parkinson's disease has been identified by a team of researchers, a finding that may one day provide a target for a new drug to prevent and potentially even cure the ... **read more »**

Neurologists Report on Promise of Statins, Estrogen, Telemedicine in Parkinson's

Apr. 28, 2014 — New approaches to understanding, treating and potentially staving off Parkinson's disease are the focus of new research findings. Studies show that factors such as estrogen exposure and statin ... **read more »**



Human Protein May Trigger Parkinson's Disease

Apr. 9, 2014 — Pathological forms of the alpha-synuclein protein present in deceased patients with Parkinson's disease are able to initiate and spread in mice and primates, researchers have found. This kind of ...

read more »